BOUTELOUA RIGIDISETA (POACEAE) NEW TO LOUISIANA

CHARLES M. ALLEN, CHRIS S. REID, and CHRIS H. DOFFITT

Department of Biology The University of Louisiana at Monroe Monroe, LA 71209, U.S.A. biallen@albha.nlu.edu

A recent grass collection from Fort Polk, in west central Louisiana, is apparently the first record for Bontelona rigidiseta (Steud.) Hitche. for the state. It is not in Allen (1993) nor Thomas and Allen (1993). In Gould (1975), the distribution is Texas, Oklahoma, and Mexico, while Smith (1994) includes this species in the Arkansas flora. The other two species of Boutelona in Louisiana are rare, with B. birsuta Lag. recorded only from Calcasieu Parish, and B. curtipendula (Michx.) Tort. from Caldwell, East Baton Rouge, Natchitoches, and Rapides parishes. The Caldwell and East Baton Rouge collections are recent, but the other three are more than 50 years old.

Voucher specimens: LOUISIANA. Vernon Parish: roadside at corner of Entrance Road and Avenue A, ca 1 mi E of La 184 on North Fort section of Fort Polk, ca 5 mi ESE of Leesville, Section 34 T2N R8W, 27 May 1999, Reid, Allen, and Doffut 423 (NLU); same locality, 25 Sep 1999, Allen 18400 (NLU).

REFERENCES

- ALLEN, C.M. 1992. Grasses of Louisiana, second edition. Cajun Prairie Habitat Pres. Soc., Eunice, LA.
- GOLLD, E.W. 1975. The grasses of Texas, Texas A. & M. Univ. Press, College Station, TX. SMITH, E.B. 1994. Keys to the flora of Arkansas. Univ. Of Arkansas Press, Fayetteville, AR.
- THOMAS, R.D. and C.M. Allen. 1993. Atlas of the vascular flora of Louisiana, Vol. 1: ferns and ferns allies, conifers, and monocoyledons. Louisiana Department of Wildlife and Fisheries, Baton Rouge, LA.